

FRONTLINE
VASCULAR CARE
2026

Abdominal Aortic Aneurysms:
Essentials and Advances in
Standard of Care




Dr. Tay Jia Sheng
MBBS (Singapore), M.Med (Surgery), FRCSed (Gen Surg), FAMS
MCR No. M11772B
Vascular, Endovascular & General Surgeon






1



2

Basics - Definition


- Aortic aneurysm
 - at least 50% increase in local aortic diameter or
 - > 30 mm absolute diameter



3

Abdominal aortic aneurysms

- Abdominal aorta aneurysms (AAA) accounts for 90% of all aneurysms
- Aortic arch, thoracic aorta and thoracoabdominal aorta is involved in ~ 10%
- 25% of AAA patients have femoral or popliteal aneurysms



4

Etiology

- Unclear – likely multifactorial process with genetic and environmental risk factors
- First-degree relatives of AAA patients have ~ 20% likelihood for development of an AAA
- 90% are thought to be due to a degenerative process
- Other causes:
 - Infection (mycotic aneurysms)
 - Connective tissue disorders
 - Arteritis

1. Marfans

2. Ehlers-Danlos

3. Loeys-Dietz

5

Epidemiology



- Aortic aneurysm usually affects elderly men
- 20th century prevalence
 - ~5% in Caucasian men aged > 65 years
 - 1.7% in women (1/3 of men)
- Recent population screening studies
 - 1.71% in men and 1.9% in women
- Trends in AAA mortality¹
 - Positively correlated with smoking and hypertension
 - Negatively correlated with obesity
- Likely due to decline in cigarette consumption and general improvement in management of cardiovascular disease risk factors

1. Ping CYM, Wu J, Tang TY, Ping HPL, Tay JB, Chikhi E. Decreases in Mortality from Abdominal Aortic Aneurysms (2001 to 2015) Is It Decreasing Even Faster? Eur J Vasc Endovasc Surg. 2021 Jun 116(1):900-907. doi: 10.1016/j.ejvs.2021.02.013. Epub 2021 Mar 24. PMID: 33773603.

6


Pathology

- Aortic aneurysmal dilatation is due to
 - reduction in medial and adventitial elastin and collagen
 - thinning of the media
 - Infiltration of lymphocytes and macrophages.
- Proposed mechanisms of aneurysm formation
 - Proteolytic degradation of aortic wall connective tissue
 - Inflammation
 - Biomechanical wall stress
 - Tangential aortic wall stress is proportional to the radius and the systemic blood pressure within the vessel (Laplace's law)
 - Increased stress and rupture risk with enlarging aneurysms
- Turbulent aortic flow due to aneurysmal morphology predisposes to thrombus formation
 - Thrombus becomes laminated against the aortic wall
 - Mural thrombus cause distal embolisation



7

Risk factors for AAA



8

Annals of Vascular Surgery

Available online 3 August 2024

In Press, Journal Pre-proof

Clinical Research

Associations Between Type 2 Diabetes Mellitus, Metabolic Traits, and Abdominal Aortic Aneurysm: A Cross-Ethnic Mendelian Randomization Analysis

Zelin Niu MD^{1,2}*, Long Cao MD, PhD^{1,3}*, Wei Guo MD, PhD¹, Hongpeng Zhang MD, PhD¹ & 

Conclusions

Type 2 diabetes mellitus protects against AAA in Europeans and East Asians. The effects of different glucose metabolism characteristics on AAA may inform AAA treatment.

9

Risk factors

- Hypertension
- Ethnicity
 - More common in Caucasians
 - Less common in Hispanics (OR, 0.7), African Americans (OR, 0.7), and Asian Americans (OR, 0.7)
- Peripheral arterial disease
- Cerebrovascular disease


10

Natural history

- Average growth rate from 0.2 to 0.3 cm per year.
- Larger AAA diameters = higher AAA growth rates.
- A wide variation between patients.
- Smoking cessation to reduce the rate of AAA growth.

11


Risk factors for rupture



12

Clinical features

- 75% Asymptomatic/Incidental
- Rupture – few present with triad of
 1. Severe abdominal or back pain
 2. Hypovolaemic shock
 3. Pulsatile abdominal mass



13

Clinical features

- Ruptured AAA should be considered in any elderly patient with unexplained hypotension / syncope and abdominal symptoms
 - Overall mortality rate of 80%
 - significant majority do not make it to hospital
 - 50% mortality rate if operated
 - Most who present have a sealed retroperitoneal haematoma with temporary haemodynamic stability.



14



AI Credit: Grok Imagine 2.0, ChatGPT Images 2.0
Original Images Credit: NYU LANGONE FEATURING THOMAS MALDONADO

15

Clinical features

- Distal embolisation of mural thrombus → Acute limb ischaemia
- Acute thrombotic occlusion of small aortic aneurysms → acute bilateral lower limb ischaemia / paralysis
- Pressure symptoms on adjacent structures
 - Dysphagia
 - Ureteric obstruction
 - Vena caval obstruction



16


Clinical features

- Inflammatory AAA may present with the triad of
 1. Low back pain
 2. Weight loss
 3. Raised erythrocyte sedimentation rate

17

Clinical features

- Rupture into an adjacent structure causing a primary aortic fistula
 - Aortocaval AVF (0.22 – 6%) with tachycardia, congestive heart failure, leg swelling, abdominal thrill, abdominal “machinery” bruit, renal failure and peripheral ischaemia.
 - 4th part of duodenum / aortoenteric fistula - herald upper gastrointestinal bleed followed by massive haemorrhage.



18

Management of AAA

- Goals
 - Prevention of rupture = asymptomatic + size criteria
 - Repair of ruptured/symptomatic AAAs
- Best medical therapy
 - Antiplatelets
 - Statins
 - BP Control
- Smoking cessation

Abdominal Aortic Aneurysms: Essentials and Advances | FVC 2026

19

AAA Annual Rupture Rate

Diameter (cm)	Annual rupture rate (%)
4 – 5.4	0.5 – 1.5
5.5 – 6	5 – 15
6 – 6.9	10 – 20
7 – 7.9	20 – 40
≥ 8	30 - 50

$T = (P \times R)$

Laplace's Law: Larger diameter = larger wall tension

Abdominal Aortic Aneurysms: Essentials and Advances | FVC 2026

20

Management of infrarenal AAA using diameter size

30 – 39mm

US Aorta
3 year
interval

M: 40 – 49 mm
F: 40 – 44 mm

US Aorta
1 year
interval

M: 50 – 54 mm
F: 45 – 49 mm

US Aorta
6 month
interval

M: ≥ 55 mm
F: ≥ 50 mm

Fit

Open or EVAR

High risk

1) Open repair or EVAR if accepts risk
2) Observe until risk of rupture > surgery

Unfit

Abdominal Aortic Aneurysms: Essentials and Advances | FVC 2026

21

Indications for Repair

Standard

- Ruptured aneurysm
 - pain due to acute expansion or imminent rupture
- Symptomatic aneurysm
 - pain due to acute expansion or imminent rupture
- Asymptomatic fusiform
 - Weigh risk of rupture vs morbidity and mortality with repair
 - Maximum AAA diameter (outer wall to outer wall)
 - Males 5.5 cm
 - Females 5.0 cm
 - Rate of growth
 - > 0.5 cm / 6 mths
 - > 1 cm / 12 mths

Glass fusiform bottle

Abdominal Aortic Aneurysms: Essentials and Advances | FVC 2026

22

Indications for Repair

Atypical

- Mycotic → Saccular aneurysm
- Penetrating aortic ulcer → Saccular aneurysm
- Recurrent aneurysm/endoleaks
- Pseudoaneurysm
- Dissection with malperfusion

Parent Artery
Neck
Dome
Sac


Abdominal Aortic Aneurysms: Essentials and Advances | FVC 2026

23


Open Repair

Abdominal Aortic Aneurysms: Essentials and Advances | FVC 2026


24



25




26




27




28




29



30




31




32




33




34



35



36



37

Outcomes

- EVAR is associated with a threefold reduction in perioperative mortality compared with propensity-matched patients undergoing elective OSR¹ including even younger patients with fewer comorbidities^{2,3}.

1. Schermerhorn ML, O'Malley AJ, Jhaveri A, Cotterill P, Pomposelli F, Landon BE. Endovascular vs. open repair of abdominal aortic aneurysms in the Medicare population. *N Engl J Med* 2008;358:464-74.

2. Siracuse JJ, Gill HL, Graham AR, Schneider DB, Connolly PH, Sedrakyan A, et al. Comparative safety of endovascular and open surgical repair of abdominal aortic aneurysms in low risk male patients. *J Vasc Surg* 2014;60:1154-8.

3. Hicks CW, Wick EC, Canner JK, Black JH 3rd, Arhuidese I, Qazi U, et al. Hospital-level factors associated with mortality after endovascular and open abdominal aortic aneurysm repair. *JAMA Surg* 2015;150:632-6.

Abdominal Aortic Aneurysms: Essentials and Advances | FVC 2020

38

Causes of poor outcomes

- Cardiovascular and pulmonary disease are leading causes of early and late death
- Despite the reduced risk of EVAR vs Open repair, EVAR has moderate to high risk of complications
- Overall EVAR outcomes are superior to those achieved with contemporary open repair, especially if done under LA/RA

Abdominal Aortic Aneurysms: Essentials and Advances | FVC 2020

39

Complications of EVAR

- Endoleaks
- Distal microembolisation – trash feet
- Endograft limb occlusion
- Access site complications e.g. pseudoaneurysms
- Graft migration / component separation
- Ischemic colitis → flexible sigmoidoscopy!
- Post-implantation syndrome
- Graft infection

Abdominal Aortic Aneurysms: Essentials and Advances | FVC 2020

40

Complications of EVAR

- Pneumonia
- Renal failure / CIN
- AMI
- Deep vein thrombosis
- Conversion to open
- Laparotomy-related complications due to open conversion
 - Adhesive bowel obstruction
 - Incisional hernia~ 10% within 6 years
 - >20% of patients treated by OSR require reoperations within 8 years
- Mandatory postoperative surveillance – annual reintervention rate 10%

Abdominal Aortic Aneurysms: Essentials and Advances | FVC 2020

41

Surveillance

- Goal of postoperative surveillance = Prevent late rupture and aneurysm-related death
 - Late rupture 8 years after EVAR is >5%
- Anastomotic aneurysm or aneurysmal dilation in the adjacent visceral aorta or iliac arteries may occur

Year	5	10	15
Incidence (%)	1	5	20

- Abdominal and pelvic CT imaging is recommend every 5 years after OSR

Abdominal Aortic Aneurysms: Essentials and Advances | FVC 2020

42

EVAR Surveillance


- CT at 1 / 6 / 12 months and yearly thereafter
- 6-month CT scan can be eliminated if the 1-month scan shows no concerning endoleak or sac enlargement.
- Further surveillance with ultrasound is safe if CT at 1 year demonstrates
 - No endoleak and stable sac size
 - Type II endoleak and a stable aneurysm size
- New endoleak, graft migration, or aneurysm sac growth >5 to 10 mm should prompt further evaluation with a CT scan

Abdominal Aortic Aneurysms: Essentials and Advances | FVC 2026

43

EVAR Surveillance

- Contrast-enhanced ultrasound (CEUS) accurate in detecting type I and type III endoleaks as well as sac enlargement
 - Eliminates ionizing radiation exposure
 - Less costly
 - Avoids risk of contrast nephropathy



Abdominal Aortic Aneurysms: Essentials and Advances | FVC 2026


44

Emergency Management Of Ruptured AAA


- Active bleeding is arrested by hypotension and tamponade by posterior parietal peritoneum
- Permissive hypotension → minimal resuscitation
- Urgent CT aortogram vs. Straight to OT
- **rEVAR** vs. Open repair
- Endovascular first approach with improving 5 year survival rates following EVAR of rAAA

Abdominal Aortic Aneurysms: Essentials and Advances | FVC 2026


45




46



47



48



49



50